MEASURE CCW: CONTRACEPTIVE CARE - ALL WOMEN AGES 15-44

Office of Population Affairs/Centers for Disease Control and Prevention

A. DESCRIPTION

The percentage of women ages 15-44 at risk of unintended pregnancy that:

- 1. Were provided a most effective or moderately effective FDA-approved method of contraception.
- 2. Were provided a long-acting reversible method of contraception (LARC).

The first measure is an intermediate outcome measure, and it is desirable to have a high percentage of women who are using the most effective or moderately effective contraceptive methods. The second measure is an access measure, and the focus is on making sure that women have access to LARC methods.

Two rates are reported for each measure, one for ages 15–20 and one for ages 21–44.

Guidance for Reporting:

- These measures apply to Medicaid enrollees ages 15–44. Two separate rates should be reported: ages 15–20 and ages 21–44. Four rates will be reported in MACPro.
- The measurement year is calendar year 2015. This specification includes ICD-9 and ICD-10 codes. In compliance with the CMS mandate to use ICD-10 codes for services provided on or after October 1, 2015, the measures should be calculated using ICD-10 codes for claims with a date of service on or after October 1, 2015.
- The National Survey of Family Growth (NSFG) can be used to interpret the results of this
 measure for the general Medicaid population. For more information, see Section E,
 "Additional Notes" and Appendix C, "How to Interpret Rates from the Performance
 Measures for Contraceptive Care."

B. DEFINITIONS

Use of a most effective method of contraception	Use of female sterilization, contraceptive implants, or intrauterine devices or systems (IUD/IUS).
Use of a moderately effective method of contraception	Use of injectables, oral pills, patch, ring, or diaphragm.
Use of a long-acting reversible method of contraception (LARC)	Use of contraceptive implants, intrauterine devices or systems (IUD/IUS).
Measurement year	Calendar year 2015.

C. ELIGIBLE POPULATION

Age	Women ages 15–44 years as of December 31 of the measurement year.
Continuous enrollment	The measurement year.

Allowable gap	No more than one gap in enrollment of up to 45 days during the continuous enrollment period. To determine continuous enrollment for a Medicaid enrollee for whom enrollment is verified monthly, the enrollee may not have more than a 1-month gap in coverage (i.e., an enrollee whose coverage lapses for 2 months [60 days] is not considered continuously enrolled).
Anchor date	December 31 of the measurement year.
Benefit	Medical or Family Planning Only Services.
Event/diagnosis	Use of contraception.

D. ADMINISTRATIVE SPECIFICATION

Denominator

Follow the steps below to define the denominator:

- Step 1 Identify all women ages 15-44 in the health plan or program.
- Step 2 Define the denominator by excluding women not at risk of unintended pregnancy because they:
 - Were infecund due to non-contraceptive reasons such as natural menopause or oophorectomy. To do this, use the codes listed in Table CCW-A.
 - Had a live birth in the last 2 months of the measurement year because there may
 not have been an opportunity to provide them with contraception. A two-month
 period was selected because the American College of Obstetricians and
 Gynecologists (ACOG) recommends having a postpartum visit by 6 weeks, and
 an additional 2 weeks was added to allow for reasonable delays in attending the
 postpartum visit. To identify live births, use the codes listed in Table CCW-D.
 - Were still pregnant at the end of the year because they were pregnant (Table CCW-B) but did not have a pregnancy outcome code indicating a non-live birth (Table CCW-C) or a live birth (Table CCW-D).

Once the exclusions are applied, the denominator includes women who were:

- Not pregnant at any point in the measurement year.
- Pregnant during the measurement year but whose pregnancy ended in the first 10 months of the measurement year, since there was adequate time to provide contraception in the postpartum period.
- Pregnant during the measurement year but whose pregnancy ended in an ectopic pregnancy, stillbirth, miscarriage, or induced abortion.

Figure CCW-A provides a flowchart for implementing these exclusion and inclusion categories.

Table CCW-A. Codes indicating sterilization for non-contraceptive reasons (i.e., hysterectomy, oophorectomy, or menopause)

ICD-9 (For dates of service from January 1, 2015 through September 30, 2015)

V49.81, V88.01, 256, 256.1, 256.2, 256.31, 256.39, 256.8, 627.1, 627.2, 627.3, 627.8, 627.9

ICD-10 (For dates of service from October 1, 2015 through December 31, 2015)

E89.40, E89.41, E28.310, E28.319, E28.39, N95.0, N95.1, N95.2, Z78.0, Z90.710, Z90.722 CPT

58150, 58152, 58180, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275,

58280, 58285, 58290, 58291, 58292, 58293, 58294, 58541, 58542, 58543, 58544, 58548,

58552, 58554, 58570, 58571, 58572, 58573, 58943, 58950, 58951, 58952, 58953, 58954,

58956, 58957, 58958, 58960

Table CCW-B. Codes indicating a pregnancy

ICD-9

V22.x, V23.x, V23.xx, V24.x, V27.x, V28.x, V28.xx, V61.6, V61.7, V72.42, V91.xx, 630-679.14

ICD-9-CM Procedure Codes

72.0-73.99, 74.0-74.20, 74.40, 74.99

ICD-10

O00-O9A, Z33.x, Z34.x, Z37.x, Z39.x

CPT

59812, 59820, 59821, 59830, 59120, 59121, 59130, 59135, 59136, 59140, 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857, 59400, 59409, 59410, 59412, 59425, 59426, 59510, 59514, 59515, 59610, 59612, 59614, 59618, 59620, 59622

Table CCW-C. Codes indicating a known miscarriage, ectopic pregnancy, stillbirth, or induced abortion

ICD-9

630-637.92, 639.0-639.9, 656.40, V27.1, V27.4, V27.7 656.41, 656.43

ICD-10

O00-O08, O36.4, Z33.2, Z37.1, Z37.4, Z37.7

CPT

59812, 59820, 59821, 59830, 59120, 59121, 59130, 59135, 59136, 59140 59840, 59841, 59850, 59851, 59852, 59855, 59856, 59857

Table CCW-D. Codes to identify a delivery resulting in a live birth

ICD-9

640.x1, 641.x1, 642.x1, 642.x2, 643.x1, 644.21, 645.x1, 646.x1, 646.x2, 647.x1, 647.x2, 648.x1, 648.x2, 649.x1, 649.x2, 650, 651.x1, 652.x1, 653.x1, 654.x1, 654.x2, 655.x1, 656.01, 656.11, 656.21, 656.31, 656.51, 656.61, 656.71, 656.81, 656.91, 657.01, 658.x1, 659.x1, 660.x1, 661.x1, 662.x1, 663.x1, 664.x1, 665.x1, 665.x2, 666.x2, 667.x2, 668.x1, 668.x2, 669.x1, 669.x2, 670.02, 671.x1, 671.x2, 672.02, 673.x1, 673.x2, 674.x1, 674.x2, 675.x1, 675.x2, 676.x1, 676.x2, 678.x1, 679.x1, 679.x2, V27.0, V27.2, V27.3, V27.5, V27.6, 670.12, 670.22, 670.32, 670.82

ICD-9-CM Procedure Codes

72.0-73.99, 74.0-74.20, 74.40, 74.99

ICD-10-CM Procedure Codes

10D00Z0, 10D00Z1, 10D00Z2, 10D07Z3, 10D07Z4, 10D07Z5, 10D07Z6, 10D07Z7, 10D07Z8, 10E0XZZ

CPT

59400, 59409, 59410, 59510, 59514, 59515, 59610, 59612, 59614, 59618, 59620, 59622

Figure CCW-A. Measure Flowchart

Step One: Identify all women ages 15-44 years

Step Two:

Define the denominator by excluding women not at risk of unintended pregnancy because they:

- Were infecund for noncontraceptive reasons
- Had a live birth in the last 2 months of the measurement year
- Were pregnant or their pregnancy outcome was unknown at the end of the measurement year

The following categories of women will remain in the denominator:

- Those who were not pregnant at any point in the measurement year
- Those who had a live birth in the first 10 months of the measurement year
- Those who had a known miscarriage, stillbirth or ectopic pregnancy, or induced abortion

Step Three:

Measure 1 Numerator. Define the numerator by identifying women who used a most or moderately effective method of contraception.

Measure 2 Numerator. Define the numerator by identifying women who used a long-acting reversible method of contraception.

Step Four:

Adjust for LARC removals and reinsertions

Step Five:

Measure 1: Calculate the rates by dividing the number who used a most or moderately effective method of contraception by the number of women in the denominator.

Measure 2: Calculate the rates by dividing the number of women who used a long-acting reversible method of contraception by the number of women in the denominator.

For both measures calculate the rates separately for:

- Adolescents (15-20 years)
- Adults (21-44 years)

Numerator for measure 1

The eligible population that used a most or moderately effective method of contraception.

- Step 3 Define the numerator by identifying women who used a most (sterilization, IUD, implant) or moderately (injection, oral pills, patch, ring, or diaphragm) effective method of contraception in the measurement year. To do this, use the codes in Table CCW-E.
- Adjust for LARC removals and re-insertions. The LARC methods can be removed at the woman's request so adjustments must be made to reflect this. Use the codes in Table CCW-F to identify women who had their IUD or implant removed at any point during the measurement year. Check to see if they had an IUD or implant reinserted on the same or a subsequent date using the codes in Table CCW-G. If there is no code indicating reinsertion, use the codes in Table CCW-E minus the Intrauterine Device (IUD/IUS) and Hormonal Implant codes to determine whether a woman was provided another most or moderately effective method in the 30 days prior to the removal (since a woman may receive a prescription for another method prior to the removal). Additionally, use all the codes in Table CCW-E to look for a subsequent most or moderately effective method in the period after the LARC removal until the end of the measurement year. If there is no code for reinsertion or provision of another most or moderately effective method, consider them as a non-user.
- Step 5 Calculate the rates by dividing the number of women who used a most or moderately effective method of contraception by the number of women in the denominator. Calculate the rates separately for adolescents and adults.

Numerator for measure 2

The eligible population that used a LARC method.

- Step 3 Define the numerator by identifying women who used a LARC in the measurement year. To do this, use the codes in Table CCW-G.
- Step 4 Adjust for LARC removals and re-insertions. The LARC methods can be removed at the woman's request so adjustments must be made to reflect this. Use the codes in Table CCW-F to identify women who had their IUD or implant removed at any point during the measurement year. Check to see if they had an IUD or implant reinserted on the same or a subsequent date through the end of the measurement year using the codes in Table CCW-G. If there is no code for reinsertion or provision of another most or moderately effective method, consider them as a non-user of LARC.
- Step 5 Calculate the rates by dividing the number of women who used a LARC method of contraception by the number of women in the denominator. Calculate the rates separately for adolescents and adults.

Table CCW-E. Codes used to identify use of most or moderately effective contraceptive methods

Description	Codes
Female	ICD-9
Sterilization	V25.2, Sterilization
	V26.51, Tubal ligation status
	ICD-9 Procedure Codes
	66.2, Procedure code bilateral endoscopic or occlusion of fallopian tubes
	ICD-10
	Z30.2, Encounter for Sterilization
	Z98.51, Tubal Ligation Status
	ICD-10 Procedure Codes
	0U574ZZ, Destruction of Bilateral Fallopian Tubes, Percutaneous Endoscopic
	Approach
	0U578ZZ, Destruction of Bilateral Fallopian Tubes, Via Natural or Artificial Opening
	Endoscopic Company of the company of
	OUL74CZ, Occlusion of Bilateral Fallopian Tubes with Extraluminal Device,
	Percutaneous Endoscopic Approach 0UL74DZ, Occlusion of Bilateral Fallopian Tubes with Intraluminal Device,
	Percutaneous Endoscopic Approach
	OUL74ZZ, Occlusion of Bilateral Fallopian Tubes, Percutaneous Endoscopic
	Approach
	0UL78DZ, Occlusion of Bilateral Fallopian Tubes with Intraluminal Device, Via
	Natural or Artificial Opening Endoscopic
	0UL78ZZ, Occlusion of Bilateral Fallopain Tubes, Via Natural or Artificial Opening
	Endoscopic
	CPT
	58600, Ligation or transection of fallopian tube(s), abdominal or vaginal approach,
	unilateral or bilateral
	58605, Ligation or transection of fallopian tube(s), abdominal or vaginal approach,
	postpartum, unilateral or bilateral, during same hospitalization (separate procedure)
	58615, Occlusion of fallopian tube(s) by device (eg, band, clip, Falope ring) vaginal
	or suprapubic approach
	58611, Ligation or transection of fallopian tube(s) when done at the time of cesarean delivery or intra- abdominal surgery (not a separate procedure) (List separately in
	addition to code for primary procedure)
	58670, Laparoscopy, surgical; with fulguration of oviducts (with or without
	transection)
	58671, Laparoscopy, surgical; with occlusion of oviducts by device (eg, band, clip, or
	Falope ring)
	58565, Hysteroscopy, surgical; with bilateral fallopian tube cannulation to induce
	occlusion by placement of permanent implants
	HCPCS
	A4264, Permanent implantable contraceptive intratubal occlusion device and delivery
	system
	- Cyclom

Description	Codes
Intrauterine	ICD-9
Device	V25.11, Encounter for insertion of intrauterine contraceptive device
(IUD/IUS)	V25.13, Encounter for removal and reinsertion of intrauterine contraceptive device
	V25.42, Surveillance of contraceptive method, intrauterine device
	V45.51, Presence of intrauterine contraceptive device
	996.32, Mechanical complication due to intrauterine contraceptive device
	IOD O Describer
	ICD-9 Procedure Codes 69.7, Insertion of intrauterine contraceptive device
	69.7, insertion of intradicinie contraceptive device
	ICD-10
	Z30.430, Encounter for insertion of intrauterine contraceptive device
	Z30.433, Encounter for removal and reinsertion of intrauterine contraceptive device
	Z30.431, Encounter for routine checking of intrauterine contraceptive device
	Z97.5, Presence of (intrauterine) contraceptive device
	T83.39XA, Other mechanical complication of intrauterine contraceptive device, initial
	encounter
	ICD 10 Presedure Codes
	ICD-10 Procedure Codes OUH97HZ, Insertion of Contraceptive Device into Uterus, Via Natural or Artificial
	Opening
	0UH98HZ, Insertion of Contraceptive Device into Uterus, Via Natural or Artificial
	Opening Endoscopic
	0UHC7HZ, Insertion of Contraceptive Device into Cervix, Via Natural or Artificial
	Opening
	0UHC8HZ, Insertion of Contraceptive Device into Cervix, Via Natural or Artificial
	Opening Endoscopic
	CPT
	58300, Insertion of IUD
	HCPCS
	J7300, Intrauterine copper contraceptive
	J7301, Levonorgestrel-releasing intrauterine contraceptive system, 13.5 mg
	J7302, Levonorgestrel- releasing intrauterine contraceptive system, 52 mg
	S4989, Contraceptive intrauterine device (e.g. progestacertiud), including implants
	and supplies
	Q0090, Levonorgestrel-releasing intrauterine contraceptive system, (skyla), 13.5 mg
	S4981, Insertion of levonorgestrel- releasing intrauterine system
	NDC
	50419042101, 50419042201, 50419042301, 51285020401

Description	Codes
Hormonal Implant	ICD-9 V25.5, Encounter for insertion of implantable subdermal contraceptive, V25.43, Surveillance of implantable subdermal contraceptive. V45.52, Presence of subdermal contraceptive implant
	CPT 11981, Insertion, non- biodegradable drug delivery implant, Implanon or Nexplanon 11983, Removal with reinsertion, non- biodegradable drug delivery implant, Implanon or Nexplanon
	HCPCS J7306, Levonorgestrel (contraceptive) implant system, including implants and supplies J7307,Etonogestrel [contraceptive] implant system, including implant and supplies
	<u>NDC</u> 00052027201, 00052027401, 00052433001
Injectable (1- month/	HCPCS J1050, Injection, medroxyprogesterone acetate
3-month)	NDC 00009074630, 00009074635, 00009470901, 00009470913, 00009737604, 00009737607, 00009737611, 00247210801, 00703680101, 00703680104, 00703681121, 23490585401, 54569370100, 54569490400, 54569552700, 54569561600, 54569621900, 54868361300, 54868410000, 54868410001, 54868525700, 55045350501, 59762453701, 59762453702, 59762453801, 59762453802, 59762453809

Description	Codes
Oral	ICD-9
Contraceptive	V25.01, Counseling and prescription of oral contraceptives
Pills	V25.41, Surveillance of contraceptive pill
5	, 3
	ICD-10
	Z30.011, Encounter for initial prescription of contraceptive pills
	Z30.41, Encounter for surveillance of contraceptive pills
	200.41, Endounter for surveinance of contradoptive pind
	HCPCS
	S4993, Contraceptive pills for birth control
	04000, Contraceptive pins for birth control
	NDC
	00008111720, 00008111730, 00008251402, 00008253505, 00008253601
	00008253605, 00052026106, 00052028306, 00052028308, 00062125100,
	00062125115, 00062125120, 00062133220, 00062141116, 00062141123, 00062171400 00062171415, 00062176100, 00062176115, 00062178100,
	00062178115, 00062179600, 00062179615, 00062190120, 00062190320,
	00062190700, 00062190715, 00062191000, 00062191015, 00093214062,
	00093209028, 00093209058, 00093313482, 00093532862, 00093542328,
	00093542358, 00093566128, 00093566158, 00093614882, 00247052028,
	00247069028, 00247069128, 00247069228, 00247139828, 00247151328,
	00247151628, 00247151728, 00247176404, 00247176421, 00247176521,
	00247198621, 00247198628, 00247200828, 00247201004, 00247201008,
	00247201028, 00247201228, 00247201328, 00247214728, 00247216928,
	00247217028, 00247223028, 00247223528, 00247226028, 00247226828,
	00378728153 00378655053, 00378727253, 00378729253, 00378730153,
	00378730853, 00430000531, 00430001005, 00430042014, 00430048214,
	00430053014, 00430053550, 00430054050, 00430057014, 00430057045,
	00430058014, 00430058045, 00430058114, 00430058514, 00430058545,
	00555034458, 00555071558, 00555900867, 00555900942, 00555901058,
	00555901258, 00555901467, 00555901658, 00555901858, 00555902058,
	00555902542, 00555902557, 00555902658, 00555902742, 00555902757,
	00555902858, 00555903270, 00555903458, 00555904358, 00555904558,
	00555904758, 00555904958, 00555905058, 00555905158, 00555905167,
	00555906458, 00555906467, 00555906558, 00555906658, 00555906667,
	00555912366, 00555913167, 00555913179, 00603359017, 00603359049,
	00603751217, 00603751249, 00603752117, 00603752149, 00603752517,
	00603752549, 00603754017, 00603754049, 00603760615, 00603760648,
	00603760715 00603760748, 00603760817, 00603760917, 00603761017,
	00603761049, 00603762517, 00603762549, 00603763417, 00603763449,
	00603764017, 00603764217, 00603766317, 00603766517, 00781405815,
	00781406015, 00781406215, 00781558307, 00781558315, 00781558336,
	00781558436, 00781558491, 00781565615, 00781565815, 16714033003,
	16714034004, 16714034604, 16714034704, 16714034804, 16714035904,
	16714036004, 16714036304, 16714036504, 16714037003, 16714040703,
	16714044004, 16714044104, 21695076928, 21695077028, 23490765301,
	23490767001, 23490769901, 24090080184, 24090096184, 35356001468,
	35356001568, 35356002168, 35356025528, 35356037028, 50102010048,
	50102012048, 50102012803, 50102013048, 50102015403, 50419040201,
	50419040203, 50419040303, 50419040503, 50419040701, 50419040703,
	50419041112, 50419041128, 50419043306, 50419043312, 50419048203,
	50419048303, 50452025115, 50458017115, 50458017615, 50458017815,
	50458019115, 50458019411, 50458019416, 50458019615, 50458019715,
	50458025115, 51285005866, 51285007997, 51285008070, 51285008198,
	51285008297, 51285008370, 51285008498, 51285008787, 51285009158,

Description	Codes
Oral	51285009287, 51285011458, 51285012058, 51285012570, 51285012698,
Contraceptive	51285012797, 51285012998, 51285013197, 51285043165, 51285054628,
Pills	51660012786, 51660057286, 52544006431, 52544014331, 52544017572,
(Continued)	52544020431, 52544021028, 52544021928, 52544022829, 52544023328,
(Continucu)	52544023528, 52544023531, 52544024531, 52544024728, 52544024828,
	52544024928, 52544025428, 52544025928, 52544025988, 52544026528,
	52544026531, 52544026829, 52544026884, 52544027428, 52544027431,
	52544027621, 52544027928, 52544029021, 52544029128, 52544029231,
	52544029241, 52544029528, 52544038328, 52544038428, 52544055028,
	52544055228, 52544055428, 52544062928, 52544063028, 52544063128,
	52544084728, 52544084828, 52544089228, 52544093628, 52544094028,
	52544094928, 52544095021, 52544095121, 52544095328, 52544095428,
	52544095931, 52544096691, 52544096728, 52544098131, 52544098231,
	54569067900, 54569068500, 54569068501, 54569068900, 54569068901,
	54569143900, 54569384400, 54569422200, 54569422201, 54569426900,
	54569427301, 54569481700, 54569487800, 54569487801, 54569489000,
	54569498400, 54569499700, 54569499800, 54569516100, 54569534900,
	54569549300, 54569549302, 54569579600, 54569579700, 54569579800,
	54569581600, 54569582600, 54569603200, 54569612800, 54569614400,
	54569627200, 54569628000, 54569628100, 54868042800, 54868044300,
	54868050200, 54868050700, 54868050801, 54868050901, 54868051600
	54868151200, 54868156400, 54868231600, 54868260600, 54868270100,
	54868377200, 54868386300, 54868394800, 54868409300, 54868423900,
	54868436900, 54868453800, 54868459000, 54868460700, 54868473000,
	54868473100, 54868474200, 54868474500, 54868475400, 54868477600,
	54868481400, 54868482800, 54868485100, 54868486000, 54868491100,
	54868502800, 54868528600, 54868532600, 54868535600, 54868582600,
	54868582800, 54868594200, 55045348506, 55045349701, 55045349801,
	55045378106, 55045378206, 55289024708, 55289088704, 55887005228,
	55887028628, 58016474701, 58016482701, 66993061128, 66993061528,
	68180084313, 68180084413, 68180084613, 68180084813, 68180085413,
	68180087611, 68180087613, 68180089213, 68180089713, 68180089813,
	68180089913, 68180090213, 68462030329, 68462030529, 68462030929,
	68462031629, 68462031829, 68462038829, 68462039429, 68462055629,
	68462056529, 68462063729, 68462064693, 00378728053, 00378728353,
	00378728753, 00378729653, 00430053750, 16714007304, 16714035903,
	16714036704, 16714040402, 16714040404, 16714040501, 16714040504,
	16714040601, 16714040604, 16714040803, 16714041304, 50419040903,
	65162031684, 65162034784 68180087513, 68180087711, 68180087713,
	68180088213, 68180088613, 68180089211, 68180089313, 75854060101
Patch	HCPCS
T dtorr	J7304, Contraceptive supply, hormone containing patch, each
	NDC
	00062192001, 00062192015, 00062192024, 50458019201, 50458019215,
	54569541300, 54868467000, 00378334053
Vaginal Ring	HCPCS
3 3	J7303, Contraceptive supply, hormone containing vaginal ring, each
	NDC NDC
	00052027301, 00052027303, 54569586500, 54868483201, 55887075401

Description	Codes
Diaphragm	<u>CPT</u>
	57170, Diaphragm or cervical cap fitting with instructions
	HCPCS
	A4266, Diaphragm for contraceptive use
	NDC
	00027013160, 00027013180, 00062330100, 00062330200, 00062330300,
	00062330400, 00062330500, 00062330600, 00062330700, 00062330800,
	00062330900, 00062331000, 00062331100, 00062331200, 00062331300,
	00062334100, 00062334200, 00062334300, 00062334400, 00062334500,
	00062334600, 00062334700, 00062334800, 00062334900, 00062335000,
	00062335100, 00062335200, 00062338100, 00062338200, 00062338300,
	00062338400, 00062338500, 00062338600, 00062338700, 00062338800,
	00062338900, 00062364103, 00062364300, 00234005100, 00234013100,
	00234013150, 00234013155, 00234013160, 00234013165, 00234013170,
	00234013175, 00234013180, 00234013185, 00234013190, 00234013195,
	00234013600, 00234013660, 00234013665, 00234013670, 00234013675,
	00234013680, 00234013685, 00234013690, 00234013695, 00396401065,
	00396401070, 00396401075, 00396401080

Table CCW-F. Codes used to identify removal/discontinued use of LARC

Description	Codes
Discontinue	ICD-9
Intrauterine	V25.12, Encounter for removal of intrauterine contraceptive device
device (IUD)	
	ICD-9 Procedure Codes
	97.71, Removal of intrauterine device
	ICD-10
	Z30.432, Encounter for removal of intrauterine contraceptive device
	ICD-10 Procedure Codes
	0UPD7HZ, Removal of Contraceptive Device from Uterus and Cervix, Via Natural or Artificial Opening
	OUPD8HZ, Removal of Contraceptive Device from Uterus and Cervix, Via Natural or Artificial Opening Endoscopic
	CPT
	58301, Encounter for removal of intrauterine contraceptive device
Discontinue	<u>CPT</u>
Implant	11976, Removal, non-biodegradable drug delivery implant, Norplant
	11982, Removal, non-biodegradable drug delivery implant, Implanon or Nexplanon

Table CCW-G. Codes used to identify use of LARC

Description	Codes
Intrauterine	ICD-9
Device (IUD/IUS)	V25.11, Encounter for insertion of intrauterine contraceptive device V25.13, Encounter for removal and reinsertion of intrauterine contraceptive device V25.42, Surveillance of contraceptive method, intrauterine device V45.51, Presence of intrauterine contraceptive device 996.32, Mechanical complication due to intrauterine contraceptive device 69.7, Insertion of intrauterine contraceptive device
	ICD-10 Z30.430, Encounter for insertion of intrauterine contraceptive device Z30.433, Encounter for removal and reinsertion of intrauterine contraceptive device Z30.431, Encounter for routine checking of intrauterine contraceptive device Z97.5, Presence of (intrauterine) contraceptive device T83.39XA, Other mechanical complication of intrauterine contraceptive device, initial encounter
	ICD-10 Procedure codes 0UH97HZ, Insertion of Contraceptive Device into Uterus, Via Natural or Artificial Opening 0UH98HZ, Insertion of Contraceptive Device into Uterus, Via Natural or Artificial Opening 0UHC7HZ, Insertion of Contraceptive Device into Cervix, Via Natural or Artificial Opening 0UHC8HZ , Insertion of Contraceptive Device into Cervix, Via Natural or Artificial Opening Endoscopic
	CPT 58300, Insertion of IUD
	HCPCS J7300, Intrauterine copper contraceptive J7301, Levonorgestrel-releasing intrauterine contraceptive system, 13.5 mg J7302, Levonorgestrel- releasing intrauterine contraceptive system, 52 mg S4989, Contraceptive intrauterine device (e.g. progestacertiud), including implants and supplies Q0090, Levonorgestrel-releasing intrauterine contraceptive system, (skyla), 13.5 mg S4981, Insertion of levonorgestrel- releasing intrauterine system
	NDC 50419042101, 50419042201, 51285020401
Hormonal Implant	ICD-9 V25.5, Encounter for insertion of implantable subdermal contraceptive, V25.43, Surveillance of implantable subdermal contraceptive. V45.52, Presence of subdermal contraceptive implant
	CPT 11981, Insertion, non- biodegradable drug delivery implant, Implanon or Nexplanon 11983, Removal with reinsertion, non- biodegradable drug delivery implant, Implanon or Nexplanon
	HCPCS J7306, Levonorgestrel (contraceptive) implant system, including implants and supplies J7307,Etonogestrel [contraceptive] implant system, including implant and supplies
	NDC 00052027201, 00052027401, 00052433001

E. ADDITIONAL NOTES

Stratification of the results by category of Medicaid eligibility (e.g., family planning waiver vs. other Medicaid eligibility) is recommended for interpretation. The National Survey of Family Growth (NSFG) can be used to interpret use of most and moderately effective contraceptive methods. NSFG should be used to interpret the results for the general Medicaid population, but the results for the family planning waiver receipients do not need to be adjusted with NSFG data. This is because the vast majority of clients who receive services from these programs are seeking contraceptive services and should therefore be considered at risk of unintended pregnancy.

The ideal denominator for a clinical performance measure of contraceptive services is all women at risk of unintended pregnancy (i.e., who are fecund, are not pregnant or seeking pregnancy, and have ever had sex). However, it is not possible to identify this population with existing claims data because there are no codes for a woman's pregnancy intention or history of sexual activity. Further, both sterilization and LARC are long-lasting but there is no systematic record of receipt of sterilization or LARC in the year(s) preceding the measurement year. These limitations can be offset by using estimates from national survey data to help interpret the measure's results and to set benchmarks that consider the limitations of claims data.

NSFG is a national survey that gathers information on family life, marriage and divorce, pregnancy, infertility, use of contraception, and men's and women's health. It is conducted by CDC's National Center for Health Statistics and generates a nationally representative sample of women and men ages 15 through 44. Approximately 5,000 individuals are interviewed each year, and updated data files are released every two years. This survey can be used to identify the portion of enrollees that are not at risk of unintended pregnancy because they never had sex, are infecund, or are trying to get pregnant. This information can then help determine the population at risk for unintended pregnancy to use as a benchmark for measure performance. For more information about the NSFG, see: http://www.cdc.gov/nchs/nsfg.htm.

See Appendix C, "How to Interpret Rates from the Performance Measures for Contraceptive Care," for examples of how to interpret performance results on this measure.